

Article Info

Accepted: 01/12/2025

1. Department of Community Medicine, Usmanu Danfodiyo University Teaching Hospital Sokoto

2. Department of Community Health, Faculty of Clinical Sciences, College of Health Science, Usmanu Danfodiyo University, PMB 2346, Sokoto, Nigeria

3. Department of Community Medicine, Federal Medical Center Gusau

*Corresponding author's email: amdanmadami@gmail.com

Cite this: *Ajoh*, 2025, vol1, i2,

FACTORS ASSOCIATED WITH VITAMIN A DEFICIENCY AMONG UNDER-FIVE CHILDREN IN URBAN AND RURAL COMMUNITIES OF NORTH-WESTERN NIGERIA.

M. D. Abdulaziz^{*1}, M. O. Oche^{1,2}, B.A Isah^{1,2}, BG Abubakar¹, MM Bello¹ Shehu A³

Abstract

Vitamin A is an essential micronutrient that is important for growth, development, immunocompetence and good vision. It is one of the micronutrients deficiencies of greatest public health concern globally, due to its high prevalence and associated health and developmental consequences. This study aimed to assess the prevalence of Vitamin A deficiency among children under five in North-Western, Nigeria.

Methods: A comparative cross-sectional study was conducted among 260 mother-child pair respondents between October 2022 and November 2022. The participants were selected by a multi-stage sampling technique. Children's serum retinol levels were determined by high-performance liquid chromatography. Data were collected using an interviewer-administered questionnaire and a data sheet. Data were processed using IBM® SPSS version 25 and analyzed using descriptive and inferential statistics.

Results: Prevalence of vitamin A was significantly higher in rural than in urban communities (33.8% vs 16.2%, $p=0.002$). In urban communities, a statistically significant association was found between vitamin A deficiency and fathers' occupation ($p < 0.001$) and vitamin A supplementation ($p = 0.017$). However, in rural areas, there was a statistically significant association between Vitamin A deficiency and fathers' occupation (<0.001), vitamin A supplement ($p =0.017$), birth order ($p= 0.010$) and Family size ($p=0.008$).

Conclusion: Vitamin A deficiency was prevalent, but a significantly higher proportion of children in the rural areas had vitamin A deficiency as compared to those in the urban areas.

Recommendation: Community health workers should educate parents about the importance of taking advantage of vitamin A supplementation and ensuring children eat foods from all food groups.

Key words: Nigeria, Vitamin A deficiency, under five children

Introduction

Hidden hunger is a term used to describe human deficiencies in essential vitamins and minerals, also known as micronutrients. Micronutrients are vitamins and minerals required in small amounts that are essential to human health, development, and growth.¹ As tiny as the amounts are, however, the consequences of their absence are severe.¹ Vitamin A is one of the most important in global public health terms; its lack represents a major threat to the health and development, particularly children and pregnant women in low and middle income countries.² Vitamin A is an essential micronutrient that is important for growth, development, immunocompetence and good vision. It is required for the differentiation and restoration of epithelial tissues in the body, as well as the normal functioning of the immune system.³ Vitamin A is one of the micronutrient deficiencies of greatest public health concern globally, due to its high prevalence and associated health and developmental consequences.⁴ Roughly 34% of children aged 6–59 months in low- and middle income countries suffer from vitamin A deficiency.^{5, 6} Vitamin A has been termed an anti-infectious agent, and it plays an important role in the visual system.² Vitamin A deficiency constitutes a major nutritional concern in the poor communities of the developing world. It is a major contributor to the morbidity and mortality of under-five children and can result in impaired resistance to infection as well as increased risk of death.³ Vitamin A deficiency impairs numerous body functions and can lead to many adverse health consequences, including xerophthalmia (dry eyes), blindness, infectious morbidity, mortality, suboptimal physical growth and anemia.² While there is overlap across such deficiencies at both population and individual levels, and the full extent of multiple deficiencies remains poorly documented, hidden hunger remains a global challenge.⁴ Understanding the epidemiology and determinants of Vitamin A deficiency within these contrasting settings is crucial for designing targeted interventions aimed at reducing the burden of deficiency and its adverse health outcomes in under-five children across Nigeria's diverse communities. This study was carried out to assess the prevalence of Vitamin A deficiency among children under five in Sokoto state, Nigeria.

MATERIALS AND METHODS

Study Design, Population, and Area

This was a comparative cross-sectional study design conducted in Sokoto. The study population comprised of mothers or caregivers with under five children in urban and rural communities of Sokoto State. Mothers or caregivers of children between 6 to 59 months of age who were residents of urban and rural communities for a minimum of twelve months were included. Mothers and caregivers who were seriously ill or having other disease conditions that made it difficult to communicate or cooperate during the interview were excluded.

Sample Size Determination

Sample size estimation was done based on the formula for the study design to compare proportions in 2 independent groups. The minimum sample size was determined using the formula.⁷

$$n = \frac{(Z_{1-\frac{\alpha}{2}} + Z_{\beta})^2 \times (P_1q_1 + p_2q_2)}{(P_1 - P_2)^2}$$

The minimum sample size of 130 subjects was recruited into the study per group (urban and rural).

Sampling technique

A multistage sampling technique was used to select respondents for the study. In stage 1, Sokoto North LGA (urban) and Wurno LGA (rural) were chosen using a simple random sampling technique by balloting. In stage 2, two wards were selected from each of the selected LGAs by a simple random sampling technique (balloting). In stage 3, one settlement was chosen from each of the four wards by a simple random sampling technique (balloting). Proportionate allocation (PA) of respondents to be enrolled in each of the settlements in each group (urban and rural) was done using the calculated sample size per group, which was 130. In stage 4, a sampling frame was obtained; systematic random sampling was used to enroll households to get respondents for the study.

Estimation of serum Vitamin A was done by the addition of ethanol to break up complexes and permit Vitamin A to partition into the heptane. The nearly colourless retinal was measured spectrophotometrically at 326 nm. The heptane extract was irradiated with UV light for about 1 hour until a steady absorbance was obtained at 326 nm. This allowed for the absorption of unwanted materials. Vitamin A was obtained by subtracting the second absorbance (A2) from the initial absorbance (A1). Since 2 ml of serum were extracted with 5 ml of heptane, the concentration of the standard equals to $37.5 \times 5/2 = 93.75$ mg/dl. Vitamin A in mg/dl equals $(A1 - A2) \times \text{concentration of standard (93.75 mg/dl)}/\text{absorbance of standard}$.⁸

Data analysis

Completed forms were downloaded from the ODK server in Excel format, data was then exported in IBM Statistical Package for Social Sciences (SPSS) version 25. Quantitative variables were summarized using the mean and standard deviation and categorical variables were summarized using frequencies and percentages. Chi-square test and Fisher's exact were used to compare sociodemographic characteristics and Vitamin A status. Independent t-test was used for comparison of mean age, weight, height and serum vitamin A. The results were presented in the form of tables and charts. Level of

significance (α) was set at 0.05, thus, any statistical test with $p < 0.05$ was considered to be statistically significant

RESULT

Table 1: Sociodemographic characteristics of the respondents

Variables	Urban (n=130) n (%)	Rural (n=130) n (%)	Test-statistics p-value
Age group of mothers (years)			
15-19	2 (1.5)	10 (7.7)	
20 – 24	13 (10.0)	22 (16.9)	
25 – 29	63 (48.5)	44 (33.8)	
30 – 34	27 (20.8)	28 (21.6)	$\chi^2= 11.059$
≥ 35	25 (19.2)	26 (20.0)	p=0.025
Mean \pm SD (years)	32.99 \pm 7.598	30.11 \pm 8.348	t=2.914 p=0.004
Religion			
Islam	125 (96.2)	130 (100)	Fischer's exact
Christianity	5 (3.8)	0	p=0.029
Tribe			
Hausa	113 (86.9)	108 (83.1)	
Fulani	9 (6.9)	7 (5.4)	
Yoruba	5 (3.8)	1 (0.8)	Fischer's exact
Igbo	3 (2.3)	0	p<0.001
Other	0	14 (10.8)	
Number of children			
1 – 4	65 (50.0)	79 (60.8)	$\chi^2= 3.051$
≥ 5	65 (50.0)	51 (39.2)	p= 0.105
Mothers' education level			
None	2 (1.5)	3 (2.3)	
Quranic only	58 (44.6)	109 (83.8)	
Primary	9 (6.9)	14 (10.8)	Fischer's exact
Secondary	45 (34.6)	3 (2.3)	P<0.001
Tertiary	16 (12.3)	1 (0.8)	
Fathers' education level			
None	0 (0.0)	0	
Quranic only	16 (12.3)	60 (46.1)	
Primary	2 (1.5)	7 (5.4)	Fischer's exact
Secondary	56 (43.1)	46 (35.4)	P<0.001
Tertiary	56 (43.1)	17 (13.1)	
Occupation of mother			
Unemployed	48 (36.9)	80 (61.5)	
Farming	0 (00.0)	0 (0.0)	Fischer's exact
Trade/business	77 (59.2)	48 (36.9)	P<0.001

Civil servant	5 (3.9)	2 (1.5)	
Occupation of father			
Unemployed	0 (0.0)	0	
Farming	6 (4.6)	39 (30.0)	$\chi^2 = 47.070$
Trade/business	66 (50.8)	74 (56.9)	p < 0.001
Civil servant	58 (44.6)	17 (13.1)	
Social class of the parent			
SC I	2 (1.5)	0	
SC II	14 (10.8)	1 (0.8)	
SC III	41 (31.5)	7 (5.4)	Fischer's exact
SC IV	63 (48.5)	58 (44.6)	P < 0.001
SC V	10 (7.7)	64 (49.2)	

χ^2 -Pearson's Chi square test; *t*- Independent *t* test

The majority of the mothers in both groups were in the age group 25-29 years. The difference in the distribution of the age groups was statistically significant ($p = 0.008$). About one third of mothers in the urban group 45 (34.6%), compared to 3 (2.3%) in the rural group, had completed their secondary education. The rural group had a higher proportion of unemployed mothers, 80 (61.5%), compared to the urban group, 48 (36.9%). There was a statistically significant difference in terms of the educational attainment and the occupation of the mothers of the children in both groups ($p < 0.001$). There was a statistically significant difference ($p < 0.001$) in the distribution of the respondents by social class, with the majority of the respondents being in social class IV and V in the Urban (48.5%) and rural groups (49.2%), respectively [Table 1].

Table 2 : Sociodemographic characteristics of children

Variables	Urban (n=130) n (%)	Rural (n=130) n (%)	Test-statistics p-value
Child's age group (months)			
12-23	32 (24.6)	28 (21.5)	$\chi^2= 0.927$ $p=0.467$
24-35	30 (23.1)	30 (23.1)	
36-47	30 (23.1)	30 (23.1)	
48-59	38 (29.2)	42 (32.3)	
Mean \pm SD (months)	34.70 \pm 14.4	35.12 \pm 14.8	
Sex			
Male	57 (43.8)	65 (50.0)	$\chi^2= 0.988$ $p=0.384$
Female	73 (56.2)	65 (50.0)	
Place of delivery			
Home	69 (53.1)	106 (81.5)	Fischer's exact $p<0.001$
Hospital	56 (43.1)	24 (18.5)	
Clinic/dispensary	4 (3.1)	0 (0.0)	
TBAs house	1 (0.8)	0 (0.0)	
Birth order			
1- 2	30 (23.0)	50 (38.5)	$\chi^2= 8.388$ $p=0.016$
3 – 4	37 (28.5)	36 (27.7)	
≥ 5	63 (48.5)	44 (33.8)	

χ^2 -Pearson's Chi-square test, t - Independent t test

The highest proportion of children in both the urban 38 (29.2%) and rural 42 (32.3%) groups were in the age group 48-59months. There was no statistically significant difference in age ($\chi^2 = 0.927$, $p = 0.467$). The mean ages in the urban and rural areas were 34.70 ± 14.4 and 35.12 ± 14.8 months, respectively ($t=0.233p=0.816$). There was no statistically significant difference in the gender distribution in the groups ($\chi^2 = 0.988$ $p = 0.384$). A higher proportion of children in the rural group, 106 (81.5%), were delivered at home compared to 69 (53.1%) in the urban group; and this was statistically significant (Fisher's exact, $p<0.001$) [Table 2].




Figure 1: Vitamin A supplement status of the children

Only a smaller proportion of children, 18 (13.8%) in urban and 14 (10.8%) rural, received all ten doses of Vitamin A (p=0.572) [Figure 2].




Figure 2: Exclusive breastfeeding status of children

In the urban areas, 23 (17.7%) of the children were exclusively breastfed compared to 14 (10.8%) of their counterparts in the rural areas. The difference was not statistically significant ($p=0.155$) [Figure 4].




Figure 3: Dietary diversity score of the family

The majority of the children, 92 (70.8%) in the rural areas, compared to 78 (60.0%) in the urban areas, had a poor dietary diversity score. The difference in proportions were not statistically significant ($p=0.090$) [Figure 5].

Table 3: Prevalence of Vitamin A deficiency among the children

Variables	Urban n (%)	Rural n (%)	Test statistics	p- value
Serum Vitamin A	n= 130	n= 130		
Normal	109 (83.8)	86 (66.2)	$\chi^2=10.851$	p=0.002
Deficient	21 (16.2)	44 (33.8)		
Mean \pm SD	35.51 \pm 16.9	28.90\pm17.1	t=3.134	p=0.001

χ^2 = Pearson's chi-square; t-independent t test

A smaller proportion, 21(16.2%) of the children in the urban areas, compared with 44 (33.8%) of those in the rural areas, have vitamin A deficiency, and this was statistically significant. (p = 0.002). The mean serum Vitamin A of children in the urban area (35.51 \pm 16.9) was more than that of the rural areas (28.90 \pm 17.1) (p= 0.001).

Table 4: Relationship between sociodemographic factors of the respondents and serum vitamin A level of the children

Variables	Urban		Test statistics and p-value	Rural		Test statistics and p-value
	Serum Vitamin A			Serum vitamin A		
	Normal	Deficient		Normal	Deficient	
Age group (months)						
12-23	26 (81.3)	6 (18.8)	$\chi^2= 0.211$	16 (57.1)	12 (42.9)	$\chi^2= 1.294$
24-59	83 (84.7)	15 (15.3)	$p= 0.782$	70 (68.6)	32 (31.4)	$p= 0.268$
Sex						
Male	49 (86.0)	8 (14.0)	$\chi^2=0.336$	38 (58.5)	27 (41.5)	$\chi^2=3.436$
Female	60 (82.2)	13 (17.8)	$p=0.636$	48 (73.8)	17 (26.2)	$p=0.095$
Tribe						
Hausa	96 (85.0)	17 (15.0)	Fisher's exact $p=0.463$	67 (62.0)	41 (38.0)	Fisher's exact $p=0.022$
Fulani	7 (77.8)	2 (22.2)		6 (85.7)	1 (14.3)	
Yoruba	4 (80.0)	1 (20.0)		1 (100.0)	0 (0.0)	
Igbo	2 (66.7)	1 (33.3)		0	0	
Others (Zabarmawa)	0	0		13 (92.9)	1 (7.1)	
Education of mother						
Nonformal	51 (85.0)	9 (15.0)	$\chi^2= 0.110$	71 (63.4)	41 (36.6)	$\chi^2=2.754$
Formal	58 (82.9)	12 (17.1)	$p=0.814$	15 (83.8)	3 (16.7)	$p= 0.114$
Occupation of mother						
Employed	44 (91.7)	4 (8.3)	$\chi^2= 1.966$	55 (69.6)	24 (30.4)	$\chi^2=1.945$
Unemployed	65 (79.3)	17 (20.7)	$p= 0.176$	31 (60.8)	20 (39.2)	$p=0.231$
Education of father						
Nonformal	15 (93.8)	1 (6.3)		36 (60.0)	24 (40.0)	

Formal	94 (82.5)	20 (17.5)	$\chi^2= 1.811$ $p= 0.467$	50 (71.4)	20 (28.6)	$\chi^2= 0.1885$ $p= 0.196$
Occupation of father						
Farming	4 (66.7)	2 (33.3)	$\chi^2=13.47$ 1 p= 0.002	13 (33.3)	26 (66.7)	$\chi^2=27.670$ p<0.001
Civil servant	42 (72.4)	16 (27.6)		12 (70.6)	5 (29.4)	
Trade/business	63 (95.5)	3 (4.5)		61 (82.4)	13 (17.6)	
Social class of parent						
Upper (SCI & II)	15 (93.8)	11 (6.2)	$\chi^2= 6.032$ $p=0.051$	1 (100.0)	0	Fisher's exact
Middle (SCIII)	36 (87.8)	5 (12.2)		6 (85.7)	1 (14.3)	$p= 0.617$
Lower (SCIV & V)	58 (79.4)	15 (20.6)		79 (64.8)	43 (35.2)	

$\chi^2=$ Pearson's chi-square

In urban areas, 6 (18.8%) of the children within the age group 12-23months had vitamin A deficiency as compared to those within 24 - 59months of age, 15 (15.3%). In urban areas, 2 (33.3%) of the children whose fathers were farmers were vitamin A deficient as compared to 16 (27.6%) and 3 (4.5%) of those whose fathers' occupations were civil service and business/trade, respectively; this was statistically significant ($p=0.002$). In the rural areas, 26 (66.7%) of children whose fathers were farmers had vitamin A deficiency compared to 5 (29.4%) and 13 (17.6%) of those whose fathers' occupations were civil service and business/trade, respectively. This was statistically significant ($p<0.001$).

Table 5: Relationship between exclusive breastfeeding, dietary diversity score, birth order, number of children, vitamin A supplement and serum vitamin A of the children

Variables	Urban		Test statistics and p-value	Rural		Test statistics and p-value
	Serum vitamin A			Serum vitamin A		
	n (%)			n (%)		
	Normal	Deficient		Normal	Deficient	
Exclusively breastfed						
Yes	18 (78.3)	5 (21.7)	Fisher's exact	9 (64.3)	5 (35.7)	Fisher's exact
No	91 (85.0)	16 (15.0)	p=0.531	77 (66.4)	39 (33.6)	p=1.000
Dietary diversity						
Good	45 (86.5)	7 (13.5)	$\chi^2=0.464$	29 (76.3)	9 (23.7)	$\chi^2=2.476$
Poor	64 (82.1)	14 (17.9)	p=0.628	57 (62.0)	35 (38.0)	p=0.154
Birth order						
1-4	60 (89.6)	7 (10.4)	$\chi^2= 2.295$	50 (58.1)	36 (41.9)	$\chi^2= 7.289$
≥ 5	49 (88.9)	14 (21.1)	p= 0.156	36 (81.8)	8 (18.2)	p= 0.010
Number of children						
1 – 4	58 (89.2)	7 (10.8)	$\chi^2=2.783$	59 (74.7)	20 (25.3)	$\chi^2=7.599$
≥ 5	51 (78.4)	14 (21.6)	p=0.152	27 (53.0)	24 (47.0)	p=0.008
Received at least 2 doses of Vitamin A supplement						
Yes	49 (92.5)	4 (7.5)	$\chi^2= 6.956$	34 (81.0)	8 (19.0)	$\chi^2= 6.069$
No	60 (77.9)	17 (22.1)	p= 0.014	52 (59.1)	36 (40.9)	p= 0.017

χ^2 = Pearson's chi-square

In the urban areas, 7(10.8%) of the children whose parents had one to four children had vitamin A deficiency compared to 14(21.6%) of those with five or more children (p =0.152). Also, 17(22.1%) of children who have not received a vitamin A supplement were vitamin A deficient as compared with 4(7.5%) of those who have received (p = 0.014). In the rural

areas, 20(25.3%) of children whose parents had one to four children were vitamin A deficient as compared to 24(47.0%) of those whose parents had five or more children ($p=0.017$). Similarly, 8(19.0%) of children who received vitamin A were vitamin A deficient compared with 36(40.9%) of those who had not received. The differences in proportions were statistically significant, $p= 0.017$ [Table 29].

Discussion

In this study, children between 48-59 months of age formed the highest proportion in both urban and rural areas, and this is similar to the findings from studies done in Imo and Bangladesh.^{9,10} A different trend was seen in a study done in Benin, Oyo and Ghana.^{11,12,13} The reason for this difference could be that the latter studies focused only on children aged 12 to 23 months compared to this study, which included children aged 12 to 59 months.

The majority of the mothers in both groups were in the age group of 25-29 years, which was similar to findings of studies carried out in Nasarawa, Oyo, Imo, Ethiopia and Iran,^{14,9,15, 13,16} in which most of the respondents fell within this age group. However, this is in contrast to a study in Ondo State of Nigeria and also in Mozambique and India^{17, 18, 19} where the majority of mothers were in the age group of 20 to 24 years. The difference obtained could be due to differences in study settings.

The prevalence of vitamin A deficiency in the present study was higher in rural (33.8%) than urban (16.2%) areas. This finding is consistent with the results of the Ghana micronutrient survey, in which vitamin A deficiency was higher in children living in rural areas (22.9%) compared to those in urban areas (18.0%).⁹ The findings in urban areas are similar to the findings in studies conducted in Osun and Saudi Arabia.^{10, 11} Similarly, the findings in the rural areas are similar to the findings in a study conducted in Sri Lanka, and another study on the micronutrient deficiencies indicators in Nigeria.^{12, 13} However, the findings of a study done in southwestern Nigeria, Egypt and the middle east contrasts these findings.^{14, 15, 16} The prevalence of vitamin A deficiency observed in this study among children in both urban and rural areas was greater than 15%, which is of public health significance.¹⁸ Thus, the importance of identifying those at risk among children and giving them Vitamin A supplements cannot be overemphasized.

A higher prevalence of vitamin A deficiency was observed among children who had not received vitamin A supplements compared to those who had received at least two doses in urban and rural areas, and the difference was statistically significant. The administration of vitamin A supplements during routine childhood immunization and Immunization Plus days might have contributed to the reduction in the prevalence of vitamin A deficiency among children who have received at least two doses of vitamin A supplementation. It was observed in this study that a small proportion of children (13.8%) in urban areas and (10.8%) in rural areas received all ten doses of Vitamin A supplementation. The Vitamin A supplementation programmes should continue and be further strengthened so as to reach all children, thereby preventing the effects of vitamin A deficiency, which are xerophthalmia (dry eyes), blindness, infectious morbidity, mortality, and suboptimal physical growth and anaemia.

The factors that were found to be significantly associated with the serum Vitamin A deficiency among the children in both groups were the occupation of the father and vitamin A supplements. In addition to the above factors, family size, birth order and tribe were found to be significantly associated with serum Vitamin A deficiency among the children in the urban group. Several studies have found some of the above factors to be significantly associated with serum Vitamin A deficiency in children.^{142, 206, 207}

Conclusion

Vitamin A was prevalent in both urban and rural areas, but a significantly higher proportion of children in the rural areas had vitamin A deficiency as compared to those in the urban areas. Community health workers should educate parents about the importance of taking advantage of childhood immunization services, including vitamin A supplementation and ensuring children eat foods from all food groups.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest

References

1. Sarvar R, Bant DD. Clinical assessment of micronutrient deficiencies among children (1- 5 years) enrolled in anganwadis of old Hubli slums, Karnataka, India. *Int J Community Med Public Health*. 2017;4(2):598-602.
2. Shemwell SA, Peratikos MB, González-Calvo L, Renom-Llonch M, Boon A, Martinho S, et al. Determinants of full vaccination status in children aged 12–23 months in Gurùé and Milange districts, Mozambique: results of a population-based cross-sectional survey. *International Health*. 2017;9(4):234-42.
3. Edem DO. Vitamin A: A review. *Asian J Clin Nutr*. 2009;1:65-82.
4. Hardin KL, Aguayo VM, Patrick Webb. Hidden hunger in South Asia: a review of recent trends and persistent challenges. *Public Health Nutrition*. 2017;21(4):785-95.
5. Stevens GA, Bennett JE, Hennocq Q. Trends and mortality effects of vitamin A deficiency in children in 138 low-income and middle-income countries between 1991 and 2013: a pooled analysis of population-based surveys. *Lancet Glob Health*3, e528–e536. 2015
6. Stevens GA, Finucane MM, De-Regil LM. Global, regional, and national trends in haemoglobin concentration and prevalence of total and severe anaemia in children and pregnant and non-pregnant women for 1995–2011: a systematic analysis of population-representative data. *Lancet Glob Health*1, e16–e25. 2013
7. Karimollah HT. "Sample size estimation in epidemiologic studies." *Caspian Journal of Internal Medicine*. 2011;2(4):289-98.
8. Valley HG, Bell M. *Practical Clinical Chemistry*. 5th ed: Oxford press; 1998.
9. Duru CB, Iwu AC, Uwakwe KA, Diwe KC, Merenu IA, Emerole CA, et al. Assessment of Immunization Status, Coverage and Determinants among under 5-Year-Old Children in Owerri, Imo State, Nigeria *Open Access Library Journal*. 2016;3(6):1-17.
10. Sarker AR, Akram R, Ali N, Sultana M. Coverage and factors associated with full immunisation among children aged 12–59 months in Bangladesh: insights from the nationwide cross-sectional demographic and health survey. *BMJ Open* 2019;9:e028020. doi:10.1136/ bmjopen-2018-028020.
11. Sally ET, Kenu E. Evaluation of access and utilization of EPI services amongst children 12-23 months in Kwahu Afram Plains, Eastern region, Ghana. *The Pan African Medical Journal*. 2017;28(238).
12. Adeleye OA, Mokogwu N. Determinants of Full Vaccination Status in a Rural Community with Accessible Vaccination Services in South-South Nigeria. *Journal of Community Medicine and Primary Health Care*. 2016;27(2):12-9.
13. Adebayo BE, Oladokun RE, Akinbami FO. Immunization Coverage in A Rural Community in Southwestern Nigeria. *J Vaccines Vaccin* (2012) 3:143. doi:10.4172/2157-7560.1000143.
14. Tabatabaei SM, Mokhtari T, Salari M, Mohammadi M. Rural-Urban Differences in Reasons for Incomplete Vaccination in Children Under Six Years, Southeast Iran 2013. *International Journal of Infection*. 2015;2(3):e28109.
15. Mekonnen AG, Bayleyegn AD, Ayele ET. Immunization coverage of 12–23 months old children and its associated factors in Minjar-Shenkora district, Ethiopia: a

- community-based study. Available from: <https://doi.org/10.1186/s12887-019-1575-7>. [last Accessed 13 April 2019].
16. Adenike OB, Adejumoke J, Olufunmi O, Ridwan O. Maternal characteristics and immunization status of children in North Central of Nigeria. *The Pan African Medical Journal* [Internet]. [cited; 26(159)].
 17. Akanbiemu FA, Fatiregun AY, Adejugbagbe AM. Nutritional status of under-fives in rural and urban communities of Southwest, Nigeria. *World Academic Journal of Community Health and Epidemiology*. 2016;2(4):064-73.
 18. Phadnis S, Khadka S, Pattanshetty S, Monteiro A. Assessment of immunization coverage among under-five year old children residing in slum settlements in an urban area in coastal Karnataka. 2015;4(5).
 19. Shemwell SA, Peratikos MB, González-Calvo L, Renom-Llonch M, Boon A, Martinho S, et al. Determinants of full vaccination status in children aged 12–23 months in Gurúé and Milange districts, Mozambique: results of a population-based cross-sectional survey. *International Health*. 2017;9(4):234-42.
 20. University of Ghana, GroundWork, University of Wisconsin-Madison, KEMRI-Wellcome Trust, UNICEF. *Ghana Micronutrient Survey 2017*. Accra, Ghana; 2017.
 21. Alquaiz AM, Gad Mohamed A, Khoja TA, Alsharif A, Shaikh S, Al Mane H, et al. Prevalence of anemia and associated factors in child bearing age women in Riyadh. *Saudi Arabia J Nutr Metab*. 2013;636585.
 22. Abolurin OO, Adegbola AJ, Oyelami OA, Adegoke SA, Bolaji OO. Vitamin A deficiency among under-five Nigerian children with diarrhoea. *Afr Health Sci* 2018;18(3):737-42
 23. Marasinghe E, Chackrewarthy S, Abeysena C, Rajindrajith S. Micronutrient status and its relationship with nutritional status in preschool children in urban Sri Lanka. *Asia Pac J Clin Nutr* 2015;24(1):144-51.
 24. Micronutrient Initiative (MI). Micronutrient Initiative. Available from: www.micronutrient.org/english/View.asp?x=596. [Last accessed 4th July 2019].
 25. Stevens GA, Bennett JE, Hennocq Q, Lu Y, de-Regil LM, Rogers L, et al. Trends and mortality effects of vitamin A deficiency in children in 138 low-income and middle-income countries between 1991 and 2013: A pooled analysis of population-based surveys. *Lancet Glob Health*. 2015; 3, e528–e36.
 26. Tawfik AA, Gohar AS, Abdel Halim S, Abdel Rahman MK. Current vitamin A status in Egypt. In *Bulletin of the National Nutrition Institute of the Arab Republic of Egypt*. National Nutrition Institute: Cairo, Egypt,. 2010:4-12.
 27. Abolurin OO, Adegbola AJ, Oyelami OA, Adegoke SA, Bolaji OO. Prevalence of Vitamin A Deficiency among Under-Five Children in South-Western Nigeria. *Niger Postgrad Med J* 2016;25:13-6.